



University of  
Zurich<sup>UZH</sup>

Zurich Open Repository and  
Archive

University of Zurich  
University Library  
Strickhofstrasse 39  
CH-8057 Zurich  
[www.zora.uzh.ch](http://www.zora.uzh.ch)

---

Year: 2018

---

## A convenient access to 1,2-diferrocenyl-substituted ethylenes via [3 + 2]-cycloelimination of 2-silylated 4,4,5,5-tetrasubstituted 1,3-dithiolanes

Mlostoń, Grzegorz ; Hamera-Faldyga, Roza ; Urbaniak, Katarzyna ; Weigand, Wolfgang ; Heimgartner, Heinz

**Abstract:** Ferrocenyl thioketones bearing a hetaryl, phenyl or alkyl group as the second substituent react with (trimethylsilyl)diazomethane at ca.  $-30^{\circ}\text{C}$  in THF solution without formation of a stable [3+2]-cycloadduct. After the spontaneous evolution of  $\text{N}_2$ , the corresponding sterically crowded 4,4,5,5-tetrasubstituted 2-silylated 1,3-dithiolanes are formed as products of the second [3 + 2]-cycloaddition of the intermediate thiocarbonyl S-methanide with the starting thioketone. After desilylation by treatment with TBAF, they are converted into the corresponding carbanions, which display different stability depending on the type of substituent. The presence of hetaryl and phenyl groups results in the exclusive formation of 1,2- diferrocenyl ethylenes. In contrast, the presence of methyl groups significantly enhances the stability of the carbanion, which by protonation yields trans-4,5-diferrocenyl-4,5-dimethyl-1,3-dithiolane.

DOI: <https://doi.org/10.1080/17415993.2018.1485920>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-152078>

Journal Article

Accepted Version

Originally published at:

Mlostoń, Grzegorz; Hamera-Faldyga, Roza; Urbaniak, Katarzyna; Weigand, Wolfgang; Heimgartner, Heinz (2018). A convenient access to 1,2-diferrocenyl-substituted ethylenes via [3 + 2]-cycloelimination of 2-silylated 4,4,5,5-tetrasubstituted 1,3-dithiolanes. *Journal of Sulfur Chemistry*, 39(5):516-524.

DOI: <https://doi.org/10.1080/17415993.2018.1485920>

## A convenient access to 1,2-diferrocenyl-substituted ethylenes via [3+2]-cycloelimination of 2-silylated 4,4,5,5-tetrasubstituted 1,3-dithiolanes

Grzegorz Mlostoń,<sup>a\*</sup> Róża Hamera-Fałdyga,<sup>a</sup> Katarzyna Urbaniak,<sup>a</sup> Wolfgang Weigand,<sup>b</sup> and Heinz Heimgartner<sup>c</sup>

<sup>a</sup>Department of Organic and Applied Chemistry, University of Łódź, Łódź, Poland;


<sup>b</sup>Institut für Anorganische und Analytische Chemie, Friedrich-Schiller-Universität Jena, Jena, Germany;

<sup>c</sup>Department of Chemistry, University of Zurich, Zurich, Switzerland

### Abstract

Ferrocenyl thioketones bearing a hetaryl, phenyl or alkyl group as the second substituent react with (trimethylsilyl)diazomethane at ca.  $-35^{\circ}\text{C}$  in THF solution without formation of a stable [3+2]-cycloadduct. After the spontaneous evolution of  $\text{N}_2$ , the corresponding sterically crowded 4,4,5,5-tetrasubstituted 2-silylated 1,3-dithiolanes are formed as products of the second [3+2]-cycloaddition of the intermediate thiocarbonyl *S*-methanide with the starting thioketone. After desilylation by treatment with TBAF, they are converted into the corresponding carbanions, which display different stability depending on the type of substituent. The presence of hetaryl and phenyl groups results in the exclusive formation of 1,2-diferrocenyl ethylenes. In contrast, the presence of methyl groups significantly enhances the stability of the carbanion, which by protonation yields *trans*-4,5-diferrocenyl-4,5-dimethyl-1,3-dithiolane.

### Graphical abstract




## Keywords

Ferrocenyl thioketones, (trimethylsilyl)diazomethane, 1,3-dithiolanes, desilylation reaction, olefination reaction

CONTACT Grzegorz Mloston [gmloston@uni.lodz.pl](mailto:gmloston@uni.lodz.pl) Department of Organic and Applied Chemistry, University of Łódź, Tamka 12, PL-91-403 Łódź, Poland

## 1. Introduction

In a series of recent publications, reactions of aryl/hetaryl thioketones **1** with diazomethane and its derivatives, such as diazoethane, 2-diazopropane and (trimethylsilyl)diazomethane, were reported [1-4]. The non-concerted pathway of the formal [3+2]-cycloaddition with participation of a diradical intermediate was demonstrated by the formation of a macrocyclic dimer of the intermediate thiocarbonyl *S*-methanide [1,5] as well as sterically crowded 4,4,5,5-tetrasubstituted 1,3-dithiolanes [2]. The latter products were formed via trapping of the intermediate diradical species by the starting thioketones in a regioselective manner. In the case of (trimethylsilyl)diazomethane **2**, 2-silylated sterically crowded 1,3-dithiolanes **3** were formed as single products (Scheme 1).




**Scheme 1.** A multi-step reaction of aryl/hetaryl thioketones **1** with TMS-diazomethane (**2**) yielding sterically crowded 4,4,5,5-tetraryl-1,3-dithiolanes **3**.

Ferrocenyl thioketones with an alkyl, aryl or hetaryl substituent are attractive building blocks for the preparation of ferrocenyl-functionalized molecules, e.g. sulfur heterocycles [6–8] and tetrasubstituted ethylenes [4,9]. The goal of present study was the reaction of ferrocenyl thioketones with (trimethylsilyl)diazomethane and subsequent desilylation of the expected 2-silylated 1,3-dithiolanes, and the observation of differences in the behavior of ferrocenyl hetaryl and alkyl ferrocenyl thioketones was of special interest.

## 2. Results and discussion


In the first part of the study, ferrocenyl hetaryl thioketones **4a–c** as well as ferrocenyl phenyl thioketone (**4d**) were tested in the reaction with TMS-diazomethane (**2**) in THF solution. The reaction mixture of equimolar amounts of both reagents was prepared at –75°C, but in contrast to aryl and hetaryl thioketones no decoloration of the solution was observed, and for that reason the mixture was slowly warmed to room temperature. Around –30°C a vigorous evolution of N<sub>2</sub> and change of the color of the solution (blue → red) were observed. The crude mixture was treated with TBAF solution at 0°C, and after 30 min, the TLC control indicated disappearance of the initially formed 1,3-dithiolane **5** (Scheme 2). The chromatographic separation delivered pure products, which were identified as (*E/Z*)-mixtures of tetrasubstituted 1,2-differocenyl ethylenes **6**. For example, in the case of the product prepared from ferrocenyl furan-2-yl thioketone (**4a**), a 10:1 mixture of isomers (<sup>1</sup>H NMR) was isolated, and after crystallization from hexane/CH<sub>2</sub>Cl<sub>2</sub> the pure major product was obtained in 54% yield. The <sup>1</sup>H NMR spectrum showed a set of signals for the ferrocenyl and furan-2-yl groups only. On the other hand, <sup>13</sup>C NMR spectra revealed a signal for the ethylene C=C unit at 129.1 ppm in addition to the characteristic signals for ferrocenyl and furan-2-yl moieties. Finally, the structure of (*E*)-**6a** was unambiguously established by X-ray crystallography [10].



**Scheme 2.** Reaction of ferrocenyl hetaryl/phenyl thioketones **4a–d** with TMS-diazomethane (**2**).

Analogous reaction courses were observed using thioketones **4b–d**. In all cases, products **6** were isolated in good yields. Whereas, **6b** and **6c** were also obtained as mixtures of (*E*)- and (*Z*)-isomers (ca. 17:1), product **6d** was isolated as a single isomer (most likely with (*E*)-configuration).


In extension of the study, TMS-diazomethane (**2**) was reacted with alkyl ferrocenyl thioketones **4e–g** under conditions presented in Scheme 3. In all experiments, the crude products obtained after desilylation were separated chromatographically, yielding two fractions. Thus, in the case of **4e**, the expected olefin **6e** formed the less polar fraction and was isolated in ca. 4% yield. Based on the <sup>1</sup>H NMR spectra, only one isomer was present, and tentatively the (*E*)-configuration can be attributed to this product. The more polar fraction contained the major product, isolated in 32% yield. The <sup>1</sup>H NMR spectrum showed the presence of the typical set of ferrocene signals along with a singlet of a Me group at 1.76 ppm and a signal of a CH<sub>2</sub> unit at 3.97 ppm. The latter is characteristic for the H<sub>2</sub>C(2) group of 4,4,5,5-tetrasubstituted 1,3-dithiolanes [11]. This hypothesis was confirmed by <sup>13</sup>C-NMR spectra, where this group was located at 27.5 ppm. Based on these data, the structure of the major product was postulated as 4,5-differocenyl-4,5-dimethyl-1,3-dithiolane (**7a**). Both products, **6e** and **7a**, were isolated as single isomers. In analogy to **6e**, the *trans*-configuration can be tentatively attributed to **7a**. The diastereoselective course of this reaction was additionally supported by the <sup>1</sup>H NMR spectrum of the crude reaction mixture containing the 2-silylated product **5e**, which indicated the presence of only one diastereoisomer.



**Scheme 3.** Reaction of alkyl ferrocenyl thioketones **4e-g** with TMS-diazomethane (**2**).

Unexpectedly, the analogous experiment performed with **4g** afforded also a mixture of products **6g** and **7c**, but in that case, the less polar alkene **6g** was the major product (37% yield) and **7c** was isolated in only ca. 5% yield. Finally, the reaction with **4f** yielded **6f** and **7b** in a ratio of 1:10.

The mechanistic interpretation of the presented results is depicted in Scheme 4. In agreement with an earlier postulated reaction sequence, the first step is the [3+2]-cycloaddition of TMS-diazomethane (**2**) with ferrocenyl thioketone **4** leading to 1,3,4-thiadiazoline **8**, which spontaneously eliminates N<sub>2</sub> generating the 1,3-diradical **9**, a mesomeric form of the thiocarbonyl *S*-methanide, acting as a reactive intermediate. This species is trapped by a second molecule of the starting thioketone **4** in a regioselective manner to give a new 1,5-diradical **10**. The subsequent 1,5-cyclization leads to 1,3-dithiolane **5** in a diastereoselective manner. Complete diastereoselectivity was observed in systems with R = phenyl or alkyl. The final step comprises desilylation of **5** affording the carbanion **11**. Apparently **its** stability depends on the type of substituents R. Whereas hetaryl groups accelerate the [3+2]-cycloelimination to give ethylenes **6**, alkyl groups enhance the stability of **11** and protonation during chromatographic workup converts this intermediate into 2-CH<sub>2</sub>-1,3-dithiolane **7**. The obtained results with derivatives bearing methyl or propyl groups suggest that the stabilizing effect of the methyl group is bigger.



**Scheme 4.** Mechanistic interpretation of the reactions of ferrocenyl thioketones **4** with TMS-diazomethane (**2**) and subsequent desilylation.

### 3. Conclusions

Ferrocenyl-substituted ethylenes attract attention as compounds of practical importance for materials and medicinal chemistry [12-new]. Some of them display interesting photo-optical properties and in the case of so called ‘ferrocifens’ their antitumor activity has extensively been studied [13-new]. The generally applied method for their synthesis is the McMurry coupling of a ferrocenyl ketone with a corresponding benzophenone derivative [14-new, 15-new]. The method presented in this study offers a new approach to 1,2-diferrocenyl ethylenes based on the application of ferrocenyl hetaryl thioketones. The obtained results show that, in comparison with diaryl and dihetaryl thioketones, ferrocenyl-containing representatives are less reactive towards (trimethylsilyl)-diazomethane and react at ca. -35°C with immediate evolution of N<sub>2</sub> to generate the corresponding thiocarbonyl *S*-methanides as reactive intermediates. The latter react with a second molecule of the starting thioketone yielding sterically crowded 4,4,5,5-tetrasubstituted 2-silylated 1,3-dithiolanes in a diastereoselective manner. These products can be easily desilylated by treatment with tetrabutylammonium fluoride to give the respective carbanion. The stability of the latter is determined by the type of the substituent of the starting ferrocenyl thioketone. In contrast to aryl and hetaryl substituents, which accelerate the [3+2]-cycloelimination leading to 1,2-diferrocenylethylenes, simple alkyl groups enhance the stability of the intermediate carbanion resulting in the subsequent protonation to give 2-CH<sub>2</sub>-1,3-dithiolane

derivatives and, therefore, they behave similar to 4,5-unsubstituted 2-silylated 1,3-dithiolanes, which are applied as masked carbanions used for C,C-bond formation [12,13].

## 4. Experimental design

### 4.1. General

All solvents were dried over appropriate drying agents and distilled before use. The  $^1\text{H}$  and  $^{13}\text{C}$  NMR spectra were measured on a Bruker Avance III instrument (600 and 150 MHz, respectively), using the solvent ( $\text{CDCl}_3$ /residual  $\text{CHCl}_3$ ) signal as reference. The IR spectra (KBr pellets) were recorded on a Nexus FT-IR spectrophotometer. The elemental analyses were determined on a Vario Micro Cube. Flash column chromatography (FCC) was carried out using Silica gel 60 (Sigma-Aldrich, 230–400 mesh). Melting points were measured in a capillary using a Stewart<sup>®</sup> SMP30. The notation Fc in this study represents ferrocenyl. Ferrocenyl thioketones **4** applied in the study were obtained by a known method according to the recently published literature protocols [3,6]. Other reagents used were commercially available chemicals.

### 4.2. Reactions of thioketones 4a–d with trimethylsilyldiazomethane (2) – a general procedure

A solution of a thioketone **4** (1 mmol) in THF (3 mL) was cooled to  $-75^\circ\text{C}$  (acetone/dry ice). Then, an equimolar amount of TMS- $\text{CHN}_2$  was added and the mixture was allowed to warm slowly to rt. Next, the solvent was evaporated, and the crude products were purified by FCC (silica gel,  $\text{CH}_2\text{Cl}_2$ /hexane 3:7).

### 4.3. Reactions of thioketones 4e–g with trimethylsilyldiazomethane (2) – a general procedure

A solution of a thioketone **4** (1 mmol) in dry THF (2 mL) was cooled to  $0^\circ\text{C}$  and the commercial solution of TMS- $\text{CHN}_2$  (2M in THF) was added in portions. The mixtures were allowed to warm to room temperature. The solvent was evaporated and the crude mixtures were dissolved in 1 mL of THF and treated with an excess of a 1M solution of TBAF in THF. After completion of the reactions, the solvent was evaporated and the crude mixtures were purified by FCC ( $\text{SiO}_2$ ,  $\text{CH}_2\text{Cl}_2$ /hexane 1:1). The main fractions



were separated on preparative plates (SiO<sub>2</sub>, CH<sub>2</sub>Cl<sub>2</sub>/hexane 3:7) to give alkenes **6e–g** (less polar fractions) and 1,3-dithiolanes **7a–c** (more polar).

#### 4.4. Desilylation of crude 1,3-dithiolane derivatives **5** – a general procedure

To the solution of crude 1,3-dithiolanes (1 mmol) in THF (4 mL), TBAF (1M, 1mL) was added at 0°C. The mixture was stirred for 20 min. Then, the solvent was evaporated and the crude product was purified by FCC (silica gel, CH<sub>2</sub>Cl<sub>2</sub>/hexane 3:7).

##### 4.4.1. 1,2-Diferrocenyl-1,2-di(furan-2-yl)ethenes (**6a**)

The product isolated after chromatography was a mixture of (*E*)/(*Z*)-isomers in a ratio of ca. 10:1. Fractional crystallization yielded the pure major component identified as the (*E*)-isomer.

(*E*)-**6a** [10]: Orange-yellow crystals; m.p. 212.0–214.0°C (from pentane/CH<sub>2</sub>Cl<sub>2</sub>); yield: 406 mg (77%). <sup>1</sup>H NMR [600 MHz, CDCl<sub>3</sub>, δ (ppm), *J* (Hz)]: 3.63–3.65 (m, 4CH(Fc)), 4.13–4.15 (m, 4CH(Fc)), 4.16 (s, 10CH(Fc)), 6.40 (d, <sup>3</sup>*J*<sub>H,H</sub> = 3.0, 2CH(Fur)), 6.54 (dd, <sup>4</sup>*J*<sub>H,H</sub> = 1.8, <sup>3</sup>*J*<sub>H,H</sub> = 3.0, 2CH(Fur)), 7.58 (brs, 2CH(Fur)). <sup>13</sup>C NMR [150 MHz, CDCl<sub>3</sub>, δ (ppm)]: 68.5, 68.9 (2 signals for 8CH(Fc)), 69.6 (10CH(Fc)), 85.4 (2C(Fc)), 109.1, 111.1, 140.8 (3 signals for 6CH(Fur)), 129.1, 153.3 (C=C, 2C(Fur)). IR (KBr): *ν* 3146w, 3109w, 3077w, 1560m, 1540m, 1495s, 1458m, 1405m, 1258s, 1236m, 1189m, 1144s, 1103s, 1049m, 1033m, 1011s, 939m, 923m, 813vs, 780m, 743vs, 596m, 527m, 476vs cm<sup>-1</sup>. ESI-MS (mixture of isomers): 528 (100, [*M*]<sup>+</sup>), 529 (50, [*M*+1]<sup>+</sup>). Anal. calcd. for C<sub>30</sub>H<sub>24</sub>Fe<sub>2</sub>O<sub>2</sub> (528.20): C 68.22, H 4.58%; found: C 68.38, H 4.61%.

(*Z*)-**6a** (minor product) was identified based on the spectra of the isolated mixture of isomers: <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>): δ 3.99 (s, 5CH(Fc)), 4.12–4.13 (m, 2CH(Fc)), 4.17–4.18 (m, 2CH(Fc)), 6.11 (d, <sup>3</sup>*J*<sub>H,H</sub> = 3.0 Hz, CH(Fur)), 6.36 (dd, <sup>4</sup>*J*<sub>H,H</sub> = 1.8 Hz, <sup>3</sup>*J*<sub>H,H</sub> = 3.0 Hz, CH(Fur)), 7.41 (brs, CH(Fur)) ppm. <sup>13</sup>C NMR (150 MHz, CDCl<sub>3</sub>): δ 67.7, 70.7 (2 signals for 4CH(Fc)), 69.3 (10CH(Fc)), 108.1, 110.6, 140.3 (3 signals for 6CH(Fur)), 129.7 (2C(Fur)) ppm.

##### 4.3.2. 1,2-Diferrocenyl-1,2-di(thiophen-2-yl)ethenes (**6b**)

The product isolated after chromatography was a mixture of (*E*)/(*Z*)-isomers in a ratio of ca. 17:1. Fractional crystallization yielded the pure major component identified as the (*E*)-isomer.

(*E*)-**6b**: Yellow-orange crystals; m.p. >158°C (from pentane/CH<sub>2</sub>Cl<sub>2</sub>); yield: 420 mg (75%). <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>): δ 3.45 (brs, 4CH(Fc)), 4.09 (brs, 4CH(Fc)), 4.14 (s, 10CH(Fc)), 6.99 (d, <sup>3</sup>J<sub>H,H</sub> = 3.0 Hz, 2CH(Thi)), 7.17 (dd, <sup>3</sup>J<sub>H,H</sub> = 3.0 Hz, <sup>3</sup>J<sub>H,H</sub> = 4.8 Hz, 2CH(Thi)), 7.48 (d, <sup>3</sup>J<sub>H,H</sub> = 4.8 Hz, 2CH(Thi)) ppm. <sup>13</sup>C NMR (150 MHz, CDCl<sub>3</sub>): δ 68.9, 69.6 (2 signals for 8CH(Fc)), 69.7 (10CH(Fc)), 86.7 (2C(Fc)), 125.3, 126.8, 127.4 (3 signals for 6 CH(Thi)), 131.0, 143.3 (C=C, 2C(Thi)) ppm. IR (KBr): ν 3086<sub>w</sub>, 2924<sub>w</sub>, 1645<sub>w</sub>, 1385<sub>w</sub>, 1250<sub>m</sub>, 1227<sub>m</sub>, 1103<sub>m</sub>, 1065<sub>m</sub>, 1039<sub>w</sub>, 999<sub>m</sub>, 925<sub>w</sub>, 837<sub>m</sub>, 814<sub>s</sub>, 723<sub>m</sub>, 695<sub>vs</sub> cm<sup>-1</sup>. ESI-MS: 560 (100, [M]<sup>+</sup>), 561 (49, [M+1]<sup>+</sup>). Anal. calcd. for C<sub>30</sub>H<sub>24</sub>Fe<sub>2</sub>S<sub>2</sub> (560.33): C 64.30, H 4.32, S 11.45, found: C 64.32, H 4.42, S 11.37.

(*Z*)-**6b** (minor product) was identified based on the spectra of a mixture of isomers: <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>): δ 3.91 (s, 10CH(Fc)), 4.17 (brs, 8CH(Fc)), 6.84–6.88 (m, 2CH(Thi)) ppm. <sup>13</sup>C NMR (150 MHz, CDCl<sub>3</sub>): δ 67.7, 70.5 (2 signals for 8CH(Fc)), 69.3 (10CH(Fc)), 87.7 (2C(Fc)), 124.0, 125.5, 126.7 (3 signals for 6CH(Thi)), 133.4, 145.6 (C=C, 2C(Thi)) ppm.

#### 4.3.3. 1,2-Diferrocenyl-1,2-di(selenophen-2-yl)ethenes (**6c**)

The product isolated after chromatography was a mixture of (*E*)/(*Z*)-isomers in a ratio of ca. 18:1. Fractional crystallization yielded the pure major component described as the (*E*)-isomer.

(*E*)-**6c**: Yellow-orange crystals; m.p. >160°C (from pentane/CH<sub>2</sub>Cl<sub>2</sub>); yield: 523 mg (80%). <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>): δ 3.66 (t, J<sub>H,H</sub> = 1.8 Hz, 4CH(Fc)), 4.11 (t, J<sub>H,H</sub> = 1.8 Hz, 4CH(Fc)), 4.14 (s, 10CH(Fc)), 7.13 (dd, <sup>4</sup>J<sub>H,H</sub> = 0.6 Hz, <sup>3</sup>J<sub>H,H</sub> = 3.6 Hz, 2CH(Sel)), 7.39 (dd, <sup>3</sup>J<sub>H,H</sub> = 3.6 Hz, <sup>3</sup>J<sub>H,H</sub> = 5.4 Hz, 2CH(Sel)), 8.16 (d, <sup>3</sup>J<sub>H,H</sub> = 5.4 Hz, 2CH(Sel)) ppm. <sup>13</sup>C NMR (150 MHz, CDCl<sub>3</sub>): δ 68.9, 70.2 (2 signals for 8CH(Fc)), 69.6 (10CH(Fc)), 86.6 (2C(Fc)), 129.2, 129.6, 131.2 (3 signals for 6CH(Sel)), 132.3, 150.4 (C=C, 2C(Sel)) ppm. IR (KBr): ν 3089<sub>m</sub>, 3044<sub>w</sub>, 2925<sub>w</sub>, 2848<sub>w</sub>, 1630<sub>w</sub>, 1446<sub>m</sub>, 1409<sub>w</sub>, 1385<sub>w</sub>, 1238<sub>m</sub>, 1213<sub>m</sub>, 1099<sub>s</sub>, 1066<sub>s</sub>, 1037<sub>m</sub>, 997<sub>m</sub>, 923<sub>m</sub>, 813<sub>vs</sub>, 765<sub>m</sub>, 723<sub>m</sub>,

686 $\nu$ s, 502 $\nu$ s  $\text{cm}^{-1}$ . ESI-MS: 654 (72,  $[M]^+$ ), 656 (100). Anal. calcd. for  $\text{C}_{30}\text{H}_{24}\text{Fe}_2\text{Se}_2$  (654.12): C 55.08, H 3.70, found: C 55.03, H 3.79.

(*Z*)-**6c** (minor product) was identified based on the spectra of a mixture of isomers:  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ ):  $\delta$  3.92 (s, 10CH(Fc)), 4.07–4.08 (m, 4CH(Fc)), 4.17–4.18 (m, 4CH(Fc)), 7.08 (d,  $^3J_{\text{H,H}} = 3.6$  Hz, 2CH(Sel)), 7.91 (d,  $^3J_{\text{H,H}} = 5.4$  Hz, CH(Sel)) ppm.  $^{13}\text{C}$  NMR (150 MHz,  $\text{CDCl}_3$ ):  $\delta$  67.7, 70.6 (2 signals for 8CH(Fc)), 69.4 (10CH(Fc)), 87.8 (2C(Fc)), 128.1, 129.0, 130.3 (3 signals for 6CH(Sel)), 134.7, 152.4 (C=C, 2C(Sel)) ppm.

#### 4.3.4. 1,2-Diferrocenyl-1,2-diphenylethene (**6d**)

The product was isolated after chromatography as a single compound described as the (*E*)-isomer.

(*E*)-**6d**: Yellow-orange crystals, m.p.  $>170^\circ\text{C}$  (decomp.); yield: 307 mg (56%).  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ ):  $\delta$  3.77 (s, 10CH(Fc)), 4.00 (brs, 4CH(Fc)), 4.14 (brs, 4CH(Fc)), 7.03–7.06 (m, 2CH<sub>arom.</sub>), 7.08–7.11 (m, 4CH<sub>arom.</sub>), 7.13–7.16 (m, 4CH<sub>arom.</sub>) ppm.  $^{13}\text{C}$  NMR (150 MHz,  $\text{CDCl}_3$ ):  $\delta$  67.6, 70.0 (2 signals for 8CH(Fc)), 69.0 (10CH(Fc)), 88.2 (2C(Fc)), 125.4, 126.9, 129.5 (3 signals for 10CH<sub>arom.</sub>), 137.4, 144.3 (C=C, 2C<sub>arom.</sub>) ppm. IR (KBr):  $\nu$  3092 $m$ , 3057 $m$ , 3023 $m$ , 2950 $m$ , 2872 $m$ , 1593 $m$ , 1577 $m$ , 1491 $m$ , 1438 $m$ , 1405 $m$ , 1381 $m$ , 1307 $m$ , 1225 $m$ , 1180 $m$ , 1107 $s$ , 1078 $m$ , 1025 $s$ , 1005 $s$ , 890 $m$ , 829 $s$ , 817 $s$ , 768 $m$ , 715 $\nu$ s, 694 $s$ , 641 $m$ , 547 $m$ , 490 $\nu$ s  $\text{cm}^{-1}$ . Anal. calcd. for  $\text{C}_{34}\text{H}_{28}\text{Fe}_2$  (548.28): C 74.48, H 5.15, found: C 74.43, H 5.28.

#### (*E*)-2,3-Diferrocenylbut-2-ene (**6e**)

The product was isolated in trace amounts (5–10 mg).  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ ):  $\delta$  2.20 (2CH<sub>3</sub>); 4.19 (s, 10CH(Fc)); 4.25–4.26 (m, 4CH(Fc)); 4.35–4.36 (m, 4CH(Fc)) ppm.  $^{13}\text{C}$  NMR (150 MHz,  $\text{CDCl}_3$ ):  $\delta$  22.6 (2CH<sub>3</sub>); 67.4, 68.9, 69.1 (3 signals for 18CH(Fc)), 90.6 (2C(Fc)), 127.8 (C=C) ppm.

#### 3,4-Diferrocenylhex-3-ene (**6f**)

The product was isolated in trace amounts (5–10 mg) as a ca. 1:1 mixture of (*E*)- and (*Z*)- isomer.  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.16 (t,  $J_{\text{H,H}} = 7.38$  Hz, 4CH<sub>2</sub>CH<sub>3</sub>), 2.64–2.69 (m, 4CH<sub>2</sub>CH<sub>3</sub>), 4.01–4.02 (m, 4CH(Fc)), 4.06 (s, 10CH(Fc)), 4.08–4.09 (m,

4CH(Fc)), 4.16 (s, 10CH(Fc)), 4.25–4.26 (m, 4CH(Fc)), 4.34–4.35 (m, 4CH(Fc)) ppm.  $^{13}\text{C}$  NMR (150 MHz,  $\text{CDCl}_3$ ):  $\delta$  15.2, 15.9 ( $2\text{CH}_2\text{CH}_3$ ), 28.8, 28.9 ( $2\text{CH}_2\text{CH}_3$ ), 66.9, 67.4, 68.8, 69.0, 69.1, 69.5 (5 signals for  $36\text{CH}(\text{Fc})$ ), 90.2, 90.8 (2 signals for  $4\text{C}(\text{Fc})$ ), 128.9, 130.9 ( $2\text{C}=\text{C}$ ) ppm.

***(E)*-4,5-Diferrocenyloct-4-ene (6g)**

Reddish crystals; mp = 111–113°C (petroleum ether); yield: 180 mg (37%).  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ ):  $\delta$  0.96 (t,  $J_{\text{H,H}} = 7.32$  Hz,  $2\text{CH}_2\text{CH}_2\text{CH}_3$ ), 1.51–1.58, 2.61–2.68 (2m,  $2\text{CH}_2\text{CH}_2\text{CH}_3$ ), 4.17 (s, 10CH(Fc)), 4.25–4.27 (m, 4CH(Fc)), 4.28–4.30 (m, 4CH(Fc)) ppm.  $^{13}\text{C}$  NMR (150 MHz,  $\text{CDCl}_3$ ):  $\delta$  14.2 ( $2\text{CH}_2\text{CH}_2\text{CH}_3$ ), 23.4, 38.4 ( $2\text{CH}_2\text{CH}_2\text{CH}_3$ ), 67.4, 69.0, 69.1 (3 signals for  $18\text{CH}(\text{Fc})$ ), 90.8 ( $2\text{C}(\text{Fc})$ ), 133.6 ( $\text{C}=\text{C}$ ). IR (KBr):  $\nu$  3094m, 2953vs, 2923m, 2867s, 1467m, 1467m, 1452m, 1258m, 1105vs, 1064m, 1041m, 1000s, 923m, 874m, 816vs, 500s, 484vs  $\text{cm}^{-1}$ . Anal. calcd. for  $\text{C}_{28}\text{H}_{32}\text{Fe}_2$  (480.12): C 70.03, H 6.72; found: C 70.02, H 6.84.

***Trans*-4,5-Diferrocenyl-4,5-dimethyl-1,3-dithiolane (7a)**

Pale yellow crystals; mp = 150–151°C (petroleum ether); yield: 160 mg (32%).  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.76 (s,  $2\text{CH}_3$ ), 3.54–3.56 (m,  $2\text{CH}(\text{Fc})$ ), 3.96 (s,  $\text{CH}_2\text{S}$ ), 4.04–4.05 (m,  $2\text{CH}(\text{Fc})$ ), 4.14 (s, 10CH(Fc)), 4.14–4.16 (m,  $2\text{CH}(\text{Fc})$ ), 4.57–4.58 (m,  $2\text{CH}(\text{Fc})$ ) ppm.  $^{13}\text{C}$  NMR (150 MHz,  $\text{CDCl}_3$ ):  $\delta$  27.5 ( $2\text{CH}_3$ ), 29.2 ( $\text{CH}_2\text{S}$ ), 66.1 ( $2\text{C}_q\text{-S}$ ), 66.9, 67.6, 68.8, 68.9, 69.1 (5 signals for  $18\text{CH}(\text{Fc})$ ), 87.5 ( $2\text{C}(\text{Fc})$ ) ppm. IR (KBr):  $\nu$  3091m, 2973m, 2931m, 1451m, 1410m, 1388m, 1359s, 1237m, 1105vs, 1058m, 1025s, 1000s, 908m, 828s, 815vs, 490vs  $\text{cm}^{-1}$ . Anal. calcd. for  $\text{C}_{25}\text{H}_{26}\text{Fe}_2\text{S}_3$  (502.29): C 59.78, H 5.22, S 12.77; found: C 59.52, H 5.50, S 12.68.

***Trans*-4,5-Diethyl-4,5-diferrocenyl-1,3-dithiolane (7b)**

Yellow crystals; mp = 147–149°C (petroleum ether); yield: 230 mg (43%).  $^1\text{H}$  NMR (600 MHz,  $\text{CDCl}_3$ ):  $\delta$  1.50 (t,  $J_{\text{H,H}} = 7.38$  Hz,  $2\text{CH}_2\text{CH}_3$ ), 1.86–1.93, 2.07–2.12 (2m,  $2\text{CH}_2\text{CH}_3$ ), 3.45–3.46 (m,  $2\text{CH}(\text{Fc})$ ), 3.79 (s,  $\text{CH}_2\text{S}$ ), 4.04–4.05 (m,  $2\text{CH}(\text{Fc})$ ), 4.08 (s, 10CH(Fc)), 4.10–4.11 (m,  $2\text{CH}(\text{Fc})$ ), 4.59–4.60 (m,  $2\text{CH}(\text{Fc})$ ) ppm.  $^{13}\text{C}$  NMR (150 MHz,  $\text{CDCl}_3$ ):  $\delta$  12.1 ( $2\text{CH}_2\text{CH}_3$ ), 29.5 ( $\text{CH}_2\text{S}$ ), 33.8 ( $2\text{CH}_2\text{CH}_3$ ), 66.1, 67.4, 68.2, 69.3, 69.4 (5 signals for  $18\text{CH}(\text{Fc})$ ), 71.5 ( $2\text{C}_q\text{-S}$ ), 88.9 ( $2\text{C}(\text{Fc})$ ) ppm. IR (KBr):  $\nu$  3095m, 2962m, 2918m, 2871m, 1447m, 1411m, 1388m, 1369m, 1262m, 1105vs, 1038m, 1029s,

998<sub>s</sub>, 934<sub>m</sub>, 816<sub>vs</sub>, 483<sub>vs</sub> cm<sup>-1</sup>. Anal. calcd. for C<sub>27</sub>H<sub>30</sub>Fe<sub>2</sub>S<sub>2</sub> (530.35): C 61.15, H 5.70, S 12.09; found: C 61.19, H 5.78, S 12.04.

***Trans-4,5-Diferrocenyl-4,5-dipropyl-1,3-dithiolane (7c)***

The product was isolated in the trace amounts (5–10 mg). <sup>1</sup>H NMR (600 MHz, CDCl<sub>3</sub>): δ 0.95 (t, *J*<sub>H,H</sub> = 7.32 Hz, 2CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 1.88–1.93 (m, 2CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 2.04–2.12, 2.13–2.21 (2m, 2CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 4.52–3.53 (m, 2CH(Fc)), 4.07–4.09 (m, 2CH(Fc)), 4.11 (s, 10CH(Fc)), 4.14–4.15 (m, 2CH(Fc)), 4.61–4.62 (m, 2CH(Fc)) ppm. <sup>13</sup>C NMR (150 MHz, CDCl<sub>3</sub>): δ 14.9 (2CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 20.2 (2CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 29.8 (CH<sub>2</sub>S), 44.1 (2CH<sub>2</sub>CH<sub>2</sub>CH<sub>3</sub>), 66.1, 67.5, 68.3, 69.1, 69.3 (5 signals for 18CH(Fc)), 70.9 (2C<sub>q</sub>-S), 89.9 (2C(Fc)) ppm.

**Acknowledgement**

Skillful performance of elemental analysis of all new compounds by Ms Agnieszka Cieślińska and Ms Hanna Jatczak (University of Łódź) is gratefully acknowledged.

**Disclosure statement**

No potential conflict of interest was reported by the authors.

**Funding**

G. M., R. H.-F. and W. W. acknowledge the Alexander von Humboldt-Stiftung (Bonn, Germany) for financial support within the ‘Institutspartnerschaft’ (Jena-Lodz, project 2018–2020).

## References

- [1] Mlostoń G, Urbaniak K, Linden A, et al. Selenophen-2-yl substituted thiocarbonyl ylides – at the borderline of dipolar and biradical reactivity. *Helv Chim Acta*. **2015**;98:453–461.
- [2] Mlostoń G, Pipiak P, Heimgartner H. Diradical reaction mechanisms in [3+2]-cycloadditions of hetaryl thioketones with alkyl or trimethylsilyl-substituted diazomethanes. *Beilstein J Org Chem*. **2016**;12:716–724.
- [3] Mlostoń G, Pipiak P, Hamera-Fałdyga R, et al. A novel application of 2-silylated 1,3-dithiolanes for the synthesis of aryl/hetaryl-substituted ethenes and dibenzofulvenes. *Beilstein J Org Chem*. **2017**;13:1900–1906.
- [4] Mlostoń G, Hamera-Fałdyga R, Heimgartner H. Silylated thiocarbonyl *S*-methanides as key intermediates in one-pot olefination reactions leading to ferrocenyl-substituted ethenes and dibenzofulvenes. *J Sulfur Chem*. **2018**; in press, DOI: 10.1080/17415993.2017.1415339
- [5] McKee ML, Mlostoń G, Urbaniak K, et al. Dimerization reactions of aryl selenophen-2-yl-substituted thiocarbonyl *S*-methanides as diradical processes: a computational study. *Beilstein J Org Chem*. **2017**;13:410–416.
- [6] Mlostoń G, Hamera R, Heimgartner H. Synthesis of ferrocenyl thioketones and their reactions with diphenyldiazomethane. *Phosphorus, Sulfur, and Silicon*. **2015**;190:2125–2133.
- [7] Kowalski K, Karpowicz R, Mlostoń G, et al. Synthesis and (spectro)electrochemistry of mixed-valent differocenyl-dihydrothiopyran derivatives. *Dalton Trans*. **2015**;44:6268–6276.
- [8] Mlostoń G, Urbaniak K, Utecht G, et al. Trifluoromethylated 2,3-dihydro-1,3,4-thiadiazoles via the regioselective [3+2]-cycloadditions of fluorinated nitrile

- imines with aryl, hetaryl, and ferrocenyl thioketones. *J Fluorine Chem.* **2016**;192:147–154.
- [9] Mlostoń G, Hamera-Fałdyga R, Jeske M, et al. Microwave-assisted reactions of  $\alpha$ -diazoketones with hetaryl and ferrocenyl thioketones. *J Sulfur Chem.* **2018**;39:47–63.
- [10] Linden A, Hamera-Fałdyga R, Mlostoń G, et al. Crystal structure of (*E*)-1,2-differocenyl-1,2-di(furan-2-yl)ethene. *Acta Cryst.* **2018**;E74:625–629.
- [11] Huisgen R, Li X, Mlostoń G, et al. Reactions of thiobenzophenone *S*-methylide with thiocarbonyl compounds. *Eur J Org Chem.* **2000**;1695–1702.
- [12] Astruc D. Why is ferrocene so exceptional? *Eur J Inorg Chem.* **2017**;6–29.
- [13] Jaouen G, Vessièrès A, Top S. Ferrocifen type anti cancer drugs. *Chem Soc Rev.* **2015**;44:8802–8817.
- [14] Jaouen G, Top S, Vessièrès A, et al. The first organometallic antioestrogens and their antiproliferative effects. *CR Acad Sci Paris, Serie IIc.* **2000**;3:89–93.
- [15] Duan X-F, Zeng J, Lü J-W, et al. Insights into the General and Efficient Cross McMurry Reactions between Ketones. *J Org Chem.* **2006**;71:9873–9876.
- [16] Degl' Innocenti A, Capperucci A, Nocentini T. 2-Trimethylsilyl-1,3-dithiolane as a masked dithiolane anion. *Tetrahedron Lett.* **2001**;27:4557–4559.
- [17] Capperucci A, Cerè V, Degl'Innocenti A, et al. Stereoselective functionalization of dithiolanes and dithianes: The first example of an axial trapping. *Synlett.* **2002**;1447–1450.